

New Fermionic Description of Quantum $S = 1/2$ Antiferromagnet

A. M. Tsvelik*

Department of Physics, Harvard University, Cambridge, Massachusetts 02138

A novel approach to $S = 1/2$ antiferromagnets with strong quantum fluctuations based on the representation of spin-1/2 operators as bylinear forms of real (Majorana) fermions is suggested. This representation has the advantage of being irreducible without any constraints on the fermionic Hilbert space. This property allowed me to write an effective theory for low-lying excitations in a spin liquid state of $S = 1/2$ antiferromagnet. It is proven that these excitations are $S = 1$ real (Majorana) fermions.

The traditional picture representing magnets as arrays of weakly interacting rigid rotors fails to describe systems with strong quantum fluctuations. Such fluctuations can be increased by a small value of spin, a frustrated interaction, or by a large value of the rank of a spin group. An alternative point of view on strongly fluctuating systems has been gradually emerging from the original Anderson's conjecture of a gapless "spin liquid" state in two-dimensional $S = 1/2$ Heisenberg systems [1]. Later he used these ideas to explain unusual magnetic properties of the copper oxide superconductors [2] which has attracted much attention. The Anderson's proposal for systems with strong fluctuations is to concentrate on bonds between spins rather than on spins themselves and thus define slow bond variables $\Delta(r, r')$ (usually called RVB - resonating valence bond - order parameter). This approach assumes that a theory of spin liquid is a lattice gauge theory.

Current efforts to develop a gauge invariant theory of spin liquid are based on the Wigner and Schwinger [3] representation of spin operators:

$$S^a = b_\alpha^\dagger \sigma_{\alpha,\beta}^a b_\beta \quad (1)$$

$$b_1^\dagger b_1 + b_2^\dagger b_2 = 2S \quad (2)$$

where σ^a are the Pauli matrices and $b^\dagger, b$ are either Bose or Fermi operators.

The above representation is used as a basis for a mean field theory where the Heisenberg exchange term is decoupled with an auxiliary field $\Delta(r, r')$ by the Hubbard-Stratonovich transformation and the constraint (2) is enforced by a local Lagrange multiplier field $\lambda(t, r)$. Depending on the expected answer some authors use the bosonic representation (see, for example, Refs.(4,5)) and some prefer the fermionic one (see Refs.(6-9)). The resulting mean field theory describes an incompressible liquid of bosons or fermions. The usual argument is that the ambiguity of statistics is irrelevant due to the incompressibility of the liquid (the constraint (2)). Meanwhile nobody has succeeded in taking the incompressibility into account. The unresolved constraint (2) reveals itself as gauge long distant forces between low-lying excitations and possibly to their confinement (see the corresponding expressions in Refs.(6-8)). The latter would mean that conventional spin wave behavior is restored. Thus the theory in the continuous limit still contains strong interactions which is unsatisfactory.

In this Letter I suggest another approach based on the special representation of spin-1/2 operators which does not contain constraints. This representation is well known in the high energy physics; the spin-1/2 operators are represented as bylinear forms of real (Majorana) fermions transforming according to the adjoint representation of the $SU(2)$ group. I begin the discussion from a brief description of the Majorana fermions.

Let us consider lattice fermions $\eta_a(r)$ ($a = 1, 2, 3$; r labels lattice sites) with the following commutation relations:

$$[\eta_a(r), \eta_b(r')]_+ = \delta_{a,b} \delta_{r,r'} \quad (3)$$

In mathematics the algebra (3) is called the Clifford algebra. For the particular case of the lattice with only one site η_a coincide with the spin-1/2 matrices. In general the commutation relations (3) can be obtained from a quantization of the following Lagrangian:

$$L_0 = \sum_r i \eta_a(r, t) \partial_t \eta_a(r, t) \quad (4)$$

It is easy to check that the following representation reproduces the commutation relations of the spin operators:

$$S^a(r) = -i\epsilon_{abc}\eta_b(r)\eta_c(r) \quad (5)$$

It follows from (3) that $\bar{S}^2 = 3/4$, i.e. the representation (5) is irreducible representation of $S = 1/2$.

The irreducibility of the representation (5) gives us an advantage of an unconstrained Hilbert space.

As the Schwinger representation (1,2) the representation (5) has a local Z_2 gauge symmetry: it is invariant with respect to the transformation

$$\eta_a(r) \rightarrow (-1)^{q(r)}\eta_a(r) \quad (6)$$

($q(r) = \pm 1$) Therefore it is 2^N -valued (N is a number of lattice sites) and the number of states of the fermions is larger than the number of spin states.

Now let us consider a Heisenberg antiferromagnet with $S = 1/2$ which Hamiltonian is given by

$$H = \sum J(r, r') S^a(r) S^a(r') - h^a S^a(r) \quad (7)$$

I use the representation (5); then the corresponding Lagrangian is $L = L_0 - H$. After a decoupling of the Heisenberg exchange interaction with an auxiliary field $\Delta(r, r')$ this Lagrangian acquires the following form:

$$L = \sum_r \left[i\eta^a(r) \partial_t \eta^a(r) + \Delta(r, r') \eta^a(r) \eta^a(r') + i h^a \epsilon_{abc} \eta^b(r) \eta^c(r) + \frac{\Delta^2(r, r')}{2J(r, r')} \right] \quad (8)$$

The Lagrangian (8) is invariant under the gauge transformation of fermions (6) and the order parameter field $\Delta(r, r')$:

$$\Delta(r, r') \rightarrow \Delta(r, r') (-1)^{[q(r) - q(r')]} \quad (9)$$

and also under transformations from the crystal lattice group:

$$\Delta \rightarrow C^{-1} \Delta C \quad (10)$$

In the mean field approximation $\Delta(r, r')$ is treated as a time-independent field determined self-consistently to get a state with minimal energy. The parameter of the expansion around the mean field saddle point is $1/n$ where $n = 3$ is a number of fermionic "colors". In principle, one can generalize this approach to the case where the spin operators are substituted by generators of the $O(n)$ symmetry group thus getting a formal parameter.

Stability of the above fixed point depends on a particular choice of $J(r, r')$. I will not discuss this question and instead concentrate on general properties of the solution provided it is stable.

In the absence of a spin long range order one can characterize the ground state by expectation values of two gauge invariant operators:

$$\epsilon(r, r') = \langle \Delta^2(r, r') \rangle \quad (11)$$

$$W(C) = \text{sign} \langle \Pi_C \Delta(r, r') \rangle \quad (12)$$

where the product in Eq.(12) is taken along a contour C .

Interesting situations arise only if a spacial variation of $|\Delta(r, r')|$ is small. Otherwise we will have trivial dimer (or spin-Peierls) states with a short range order. Neglecting fluctuations of $|\Delta|$ and substituting it for a coordinate independent quantity Δ_0 found from the condition of minimal free energy we get from Eq.(8) the following effective Lagrangian for low energy modes of the theory:

$$L = \sum_r i\eta_a(r) \partial_t \eta_a(r) + i h^a \epsilon_{abc} \eta^b(r) \eta^c(r) + \sum_{\langle r, r' \rangle} i \Delta_0 \eta_a(r) \sigma(r, r') \eta_a(r') \quad (13)$$

where $\sigma(r, r') = \pm 1$ is a Z_2 gauge field defined on the links of the lattice with nonvanishing $J(r, r')$.

The Lagrangian (13) allows a further simplification if in the ground state $W(C) = 1$ for all loops (uniform states). The uniform state could be stable provided there are no flat parts on the Fermi surface. Usually such parts are removed by interactions with next nearest neighbours.

For the further analysis it is convenient to make a Fourier transformation of the fermionic fields. From Eq.(3) it follows that the transformed fields satisfy the following commutation relations:

$$[\eta^a(\vec{k}), \eta^b(\vec{k}')] = \delta_{ab} \delta_{\vec{k} + \vec{k}', 0} \quad (14)$$

Then from Eq.(8) I derive the Hamiltonian for the uniform state (the Zeeman term is omitted):

$$H = \sum_{\vec{k}} \epsilon(\vec{k}) \eta^a(-\vec{k}) \eta^a(\vec{k}) + \sum_{\vec{q}, \vec{p}, \vec{k}} J(\vec{p} - \vec{k}) : \eta^a(\vec{k}) \eta^a(-\vec{k} + \vec{q}) :: \eta^b(-\vec{p}) \eta^b(\vec{p} - \vec{q}) : \quad (15)$$

where the dots denote the normal ordering and

$$\epsilon(\vec{k}) = \langle \eta^a(-\vec{k}) \eta^a(\vec{k}) \rangle = \sum_{\langle r, r' \rangle} J(r, r') \cos[\vec{k}(\vec{r} - \vec{r}')] \quad (16)$$

The energy $\epsilon(\vec{k})$ vanishes on some surface; since according to Eq.(16) $\epsilon(\vec{k}) = -\epsilon(-\vec{k})$ this surface divides the Brillouin zone into two equal parts Ω_+ (where $\epsilon > 0$) and Ω_- ($\epsilon < 0$). Then $\eta(\vec{k})$ and $\eta(-\vec{k})$ with $\vec{k}$ belonging to Ω_- are creation and annihilation operators which returns us to the conventional Fermi liquid picture. Fermions fill a half of the Brillouin zone which guarantees that the spin sum rules are satisfied. In fact, the uniform state is a chargeless Fermi liquid. It follows from the fact that due to a nonsingular character of the interaction in Eq.(15) all scattering effects become irrelevant close to the Fermi surface. Therefore fermions with small energies propagate coherently. Thus this state of a quantum antiferromagnet does have fermions as elementary excitations, but these fermions have spin 1 instead of the usual 1/2 since they belong to the adjoint representation of the SU(2) group! As in a conventional Fermi liquid the system has a linear specific heat $C_v \propto T$; the magnetic susceptibility $\chi(\omega, \vec{q})$ is almost $\vec{q}$ -independent at small $\vec{q}$, has a weak singularity at $|\vec{q}| = 2p_F$ and exhibits a significant imaginary part in the broad range of frequencies $\omega \sim J$ without an energy gap. Such behavior has been observed in various strongly fluctuated magnetic systems.

Except of the uniform states another possibility exists which arises when the mean field ground state favors $W(C) = -1$ for certain loops. Such ground states has been intensively discussed in the literature as flux (or chiral) states (see Refs.(6-10)). The present case differs from the conventional formulation of the problem in two aspects: the effective hopping integrals $\Delta(r, r')$ are now real and the fermionic filling factor is always equal to 1/2.

A flux state is periodic, but its periodicity does not coincide with the initial periodicity of the lattice. A new elementary cell arises which is defined as a minimal cell surrounded by contours with $W(C) = 1$. Since the initial symmetry is broken a flux phase can arise at low temperatures only via a phase transition. Apparently in the high temperature phase the fermions are confined into ordinary spins and the transition release them. Since the broken symmetry is discrete, such transition may happen even in two dimensions. Whether it happen or not is not clear, however. For the lattice Z_2 gauge theory which is similar to the theory I discuss, it was established in Ref.11 that there is no phase transition in $d = 2$ and it occurs only in $d = 3$ as a second order transition. Therefore the problem of existence of two dimensional flux phase of a quantum antiferromagnet requires further studies. As far as hypothetical three dimensional flux states are concerned they would have the following properties. As we know, the spectrum of fermions in a flux state is either gapful or has conical singularities: $\epsilon(\vec{k}) \propto |\vec{k} - \vec{k}_0|$ [10]. The specific heat in the latter case looks like for an ordinary antiferromagnet: $C_v \propto T^3$, but the magnetic properties are essentially different. In particular, the magnetic susceptibility behaves as $\chi \propto h^2$. It cannot be excluded that the remaining interactions can cause singularities in the nonlinear magnetic susceptibility. I emphasise that elementary excitations are again $S = 1$ fermions and not spinons as it follows from the ground state wave function postulated in Ref.10.

Unfortunately, I have not managed to generalize the described approach to the $SU(2/1)$ group which is necessary if one wants to include effects of doping. It prevents me from a straightforward application of this technique to the Hubbard or the $t - J$ models. At the present time the described procedure can be used only for problems with well defined spin operators, for example, for the Kondo lattice problem [12].

I am grateful to P. W. Anderson and P. Coleman for intensive and valuable discussions and to P. B. Wiegmann for sending me his preprint. This work was supported by the National Science Grant Foundation, through grant DMR 91-15491 and through the Harvard University Materials Research Laboratory.

* The permanent address is Department of Physics, University of Oxford, 1 Keble Road, Oxford OX1 3NP, England.

[1] P. W. Anderson, Mater. Res. Bull. **8**, 153 (1973).

[2] P. W. Anderson, Science **235**, 1196 (1987).

[3] J. Schwinger, US Atomic Energy Commission, NYO-3071 (1952), unpublished report; E. Wigner, private communication.

[4] D. P. Arovas and A. Auerbach, Phys. Rev. **B 38**, 316 (1988); Phys. Rev. Lett. **61**, 617 (1988).

- [5] P. Chandra, P. Coleman and A. I. Larkin, J. Phys. Condens. Matt. **2**, 7933 (1990).
- [6] D. Khvestchnko and P. B. Wiegmann, Mod. Phys. Lett **3**, 1383 (1989); ibid. **4**, 17 (1990).
- [7] P. B. Wiegmann, unpublished.
- [8] X.-G. Wen, F. Wilczek and A. Zee, Phys. Rev. **B39**, 11413 (1989).
- [9] R. B. Laughlin and Z. Zou, Phys. Rev. **B41**, 664 (1989).
- [10] S. B. Libby, Z. Zou and R. B. Laughlin, Nucl.Phys. **B348**, 693 (1991).
- [11] R. Balian, J. M. Drouffe and C. Itzykson, Phys. Rev. **D11**, 2098 (1975).
- [12] P. Coleman, F. Miranda and A. M. Tsvelik, in preparation.